

Bexsero vaccine error warning

**HIGH
ERROR
RATE**

Additional checks are recommended due to a high number of spacing errors that have occurred.

As well as all the normal checks, we strongly recommend the use of the following to double check you have the correct spacing and schedule:

- **Confirm details including current age**
- **Double check past vaccine history.** For PMS, check Immunisation Schedule, and All Immunisation tabs for Medtech.) On AIR, check number of MenB doses, age when given and spacing between doses. Review Well Child book if available.

Name		Age when Bexsero course started:	
Date given	Dose	Space between doses	
	1		
	2		
	B*		
Are you sure the vaccine is due today?			
If you give the vaccine today, will the spacing between doses be correct?			
Is a booster needed? *Only required for those starting vaccine at 2 years or under.			
If additional doses are required, date when due? Arrange appointment if possible			
Has this been double checked ?			

BEXSERO VACCINATION SCHEDULES

Age when Bexsero first dose given	Number and timing of doses
Infants >6 weeks to <11 months	<ul style="list-style-type: none"> - 2 doses separated by 8 weeks - Plus booster dose aged ≥12 months or a minimum of 6 months after second dose, whichever is later
Infants ≥12 to ≤23 months (for first dose)	<ul style="list-style-type: none"> - 2 doses separated by 8 weeks - Plus booster dose given 12-24 months after second dose
Children ≥24 months (for first dose) and adults*	<ul style="list-style-type: none"> - 2 doses separated by 8 weeks - Booster doses are funded for high-risk groups

Please refer to the **7 Rights of Bexsero vaccine administration prompt card** for more information [here](https://tinyurl.com/3zatfxp7) (tinyurl.com/3zatfxp7)

Call 0800 IMMUNE (0800 466 863) for clinical advice